

Adam Seidl, MD
Assistant Professor
Shoulder/Elbow Surgery
Department of Orthopedics
University of Colorado



Lone Tree Health Center
(720) 848-2200

Anschutz Medical Campus
(720) 848-1900

Rehabilitation Protocol Anterior Instability Repair

General Instructions:

- Therapy is to begin 2 weeks after the surgery.
- The patient should work with their therapist 1-3X per week until released by the surgeon.
- Do not add or skip any part of this program. If you have concerns please contact my office.
- Goals: The 2 main goals of this physical therapy program are to:
 - Have full active and passive range of motion by 3 months after surgery
 - Return to sport by 18-24 weeks after surgery

Sling Wear:

- Patients must wear their sling **at all times** for the first 6 weeks after surgery, this does include while they are sleeping. Patients may only remove the sling to perform therapy exercises and for showering.

Ice:

- The use of ice or ice machine is encouraged to help control pain and inflammation after surgery.

Questions/Concerns:

- Therapists – if you have questions or concerns, please contact me directly or my office. Email is best for non-urgent issues: Adam.Seidl@ucdenver.edu

PROTOCOL

All time points are based on time since surgery:

0-2 Weeks

- Wrist and Elbow ROM Only

Adam Seidl, MD
Assistant Professor
Shoulder/Elbow Surgery
Department of Orthopedics
University of Colorado



Lone Tree Health Center
(720) 848-2200

Anschutz Medical Campus
(720) 848-1900



2 Weeks

- Passive/active assist forward elevation (FE) to 90 degrees
- Passive/active assist external rotation (ER) to neutral with arm at the side
- Gentle Isometrics (no ER/IR)



4 Weeks

- Passive/active assist forward elevation (FE) to 120 degrees
- Passive/active assist external rotation (ER) to 20 degrees with arm at the side
- Passive/active assist Abduction to 90 degrees
- Scapula protraction/retraction (with arm in sling until 6 weeks post-op)
- No combined Abduction and ER!



6 Weeks

- May discontinue sling usage, unless in crowd, or on slippery surfaces
- Unlimited passive/active assist in FE
- May begin active motion in all planes
- Posterior glides OK (no anterior)
- Resisted isometrics (no IR)

Adam Seidl, MD
Assistant Professor
Shoulder/Elbow Surgery
Department of Orthopedics
University of Colorado



Lone Tree Health Center
(720) 848-2200

Anschutz Medical Campus
(720) 848-1900

8 Weeks

- Continue to progress with active motion
- May slowly progress to resisted exercise with therabands



12 Weeks

- May begin sport specific exercises



18-24 Weeks

- Return to play with approval of surgeon