

Adam Seidl, MD
Associate Professor
Shoulder/Elbow Surgery
Department of Orthopedics
University of Colorado



Orthopedics
UNIVERSITY OF COLORADO

Highlands Ranch Hospital
(720) 516-4090
Inverness
(303) 694-3333
Anschutz Medical Campus
(720) 848-1900

Rehabilitation Protocol Posterior Instability Repair

General Instructions:

- Therapy is to begin 2 weeks after the surgery.
- The patient should work with their therapist 1-3X per week until released by the surgeon.
- Do not add or skip any part of this program. If you have concerns please contact my office.
- Goals: The 2 main goals of this physical therapy program are to:
 - Have full active and passive range of motion by 3 months after surgery
 - Return to sport by 18-24 weeks after surgery

Sling Wear:

- Patients must wear their sling **at all times** for the first 6 weeks after surgery, this does include while they are sleeping. Patients may only remove the sling to perform therapy exercises and for showering.

Ice:

- The use of ice or ice machine is encouraged to help control pain and inflammation after surgery.

Questions/Concerns:

- Therapists - if you have questions or concerns, please contact me directly or my office. Email is best for non-urgent issues: Adam.Seidl@ucdenver.edu

PROTOCOL

All time points are based on time since surgery:

0-2 Weeks

- Wrist and Elbow ROM Only

Adam Seidl, MD
Associate Professor
Shoulder/Elbow Surgery
Department of Orthopedics
University of Colorado



Orthopedics
UNIVERSITY OF COLORADO

Highlands Ranch Hospital
(720) 516-4090
Inverness
(303) 694-3333
Anschutz Medical Campus
(720) 848-1900

2 Weeks

- Passive/active assist forward elevation (FE) to 90 degrees



4 Weeks

- Passive/active assist forward elevation (FE) to 120 degrees




- Passive/active assist Abduction to 90 degrees



- Isometrics - no External Rotation (ER)
- No combined Abduction and IR!

6 Weeks

- May discontinue sling usage, unless in crowd, or on slippery surfaces
- Unlimited passive/active assist in FE
- May begin active motion in all planes - to remain less than passive limits
- Resisted isometrics (no ER)
- PRE's - No ER/IR
- Scapular stabilizers - protraction/retraction



- Anterior glides OK if needed - no posterior

Adam Seidl, MD
Associate Professor
Shoulder/Elbow Surgery
Department of Orthopedics
University of Colorado



Orthopedics
UNIVERSITY OF COLORADO

Highlands Ranch Hospital
(720) 516-4090
Inverness
(303) 694-3333
Anschutz Medical Campus
(720) 848-1900

8 Weeks

- Passive/active assist internal rotation to 30 degrees with arm at the side
- Passive/active assist internal rotation at 45 degrees abduction to 30 degrees
- Continue to progress with active motion
- May slowly progress to resisted exercise with therabands



10 Weeks

- Unlimited internal rotation - passive and active
- PRE's - with IR/ER

14 Weeks

- May begin sport specific exercises



- Posterior glides OK if needed

20-24 Weeks

- Return to play with approval of surgeon