

**Adam Seidl, MD**  
Assistant Professor  
Shoulder/Elbow Surgery  
Department of Orthopedics  
University of Colorado



**Orthopedics**  
UNIVERSITY OF COLORADO

**Highlands Ranch Hospital**  
(720) 516-4090  
**Inverness**  
(303) 694-3333  
**Anschutz Medical Campus**  
(720) 848-1900

## Rehabilitation Protocol Posterior Instability Repair

### **General Instructions:**

- Therapy is to begin 2 weeks after the surgery.
- The patient should work with their therapist 1-3X per week until released by the surgeon.
- Do not add or skip any part of this program. If you have concerns please contact my office.
- Goals: The 2 main goals of this physical therapy program are to:
  - Have full active and passive range of motion by 3 months after surgery
  - Return to sport by 18-24 weeks after surgery

### **Sling Wear:**

- Patients must wear their sling **at all times** for the first 6 weeks after surgery, this does include while they are sleeping. Patients may only remove the sling to perform therapy exercises and for showering.

### **Ice:**

- The use of ice or ice machine is encouraged to help control pain and inflammation after surgery.

### **Questions/Concerns:**

- Therapists – if you have questions or concerns, please contact me directly or my office. Email is best for non-urgent issues: [Adam.Seidl@ucdenver.edu](mailto:Adam.Seidl@ucdenver.edu)

### **PROTOCOL**

**All time points are based on time since surgery:**

#### **0-2 Weeks**

- Wrist and Elbow ROM Only

**Adam Seidl, MD**  
Assistant Professor  
Shoulder/Elbow Surgery  
Department of Orthopedics  
University of Colorado



**Orthopedics**  
UNIVERSITY OF COLORADO

**Highlands Ranch Hospital**  
(720) 516-4090  
**Inverness**  
(303) 694-3333  
**Anschutz Medical Campus**  
(720) 848-1900


**2 Weeks**

- Passive/active assist forward elevation (FE) to 90 degrees



**4 Weeks**

- Passive/active assist forward elevation (FE) to 120 degrees




- Passive/active assist Abduction to 90 degrees



- Isometrics – no External Rotation (ER)
- No combined Abduction and IR!

**6 Weeks**

- May discontinue sling usage, unless in crowd, or on slippery surfaces
- Unlimited passive/active assist in FE
- May begin active motion in all planes – to remain less than passive limits
- Resisted isometrics (no ER)
- PRE's – No ER/IR
- Scapular stabilizers – protraction/retraction



- Anterior glides OK if needed – no posterior

**Adam Seidl, MD**  
Assistant Professor  
Shoulder/Elbow Surgery  
Department of Orthopedics  
University of Colorado



**Orthopedics**  
UNIVERSITY OF COLORADO

**Highlands Ranch Hospital**  
(720) 516-4090  
**Inverness**  
(303) 694-3333  
**Anschutz Medical Campus**  
(720) 848-1900

### **8 Weeks**

- Passive/active assist internal rotation to 30 degrees with arm at the side
- Passive/active assist internal rotation at 45 degrees abduction to 30 degrees
- Continue to progress with active motion
- May slowly progress to resisted exercise with therabands



### **10 Weeks**

- Unlimited internal rotation – passive and active
- PRE's – with IR/ER

### **14 Weeks**

- May begin sport specific exercises



- Posterior glides OK if needed

### **20-24 Weeks**

- Return to play with approval of surgeon