

Adam Seidl, MD
Assistant Professor
Shoulder/Elbow Surgery
Department of Orthopedics
University of Colorado



Orthopedics
UNIVERSITY OF COLORADO

Highlands Ranch Hospital
(720) 516-4090
Inverness
(303) 694-3333
Anschutz Medical Campus
(720) 848-1900

Rehabilitation Protocol SLAP Repair

Phase I: Protect repair, restore ROM (*0 to 6 weeks*)

- May remove dressing and shower postop day # 3.
- Sutures are all underneath the skin and will dissolve on their own.
- Ice or cold flow systems encouraged for the first week at a minimum: should be used 3-4 times per day.
- Sling should be in place when not performing exercises.
- Scapular ROM exercises.
- No lifting with involved extremity.
- Avoid pure active shoulder exercises, resisted biceps exercises.
- Avoid shoulder rotation ROM from abducted position and cross body motions.
- Initiate exercise program 3 times per day:
 - Immediate elbow, forearm and hand range of motion out of sling
 - Pendulum exercises
 - Passive and active assistive ROM – flexion and scapular plane elevation to tolerance, ER to 45 (pulleys, supine gravity assisted exercises)
 - Emphasize home program

PHASE II: Advance ROM (*6 to 12 weeks*)

- May discontinue sling.
- Lifting restriction of 10 pounds should be reinforced with patient until 3 months.
- Advance to AROM exercises.
- Advance PROM and capsular mobility as tolerated.
- Avoid rotational ROM in abducted position until 8 weeks postop.
- Initiate rotator cuff and gentle biceps strengthening (up to 10 lbs).

Phase III: Restore function (*> 3 months*)

- Discontinue lifting restriction.
- Advance to higher-level rotator cuff and scapular stabilizer strengthening.
- Start throwing progression if applicable.